

COVID-19 is an emerging, rapidly evolving situation.

Get the latest public health information from CDC: <https://www.coronavirus.gov>.

Get the latest research from NIH: <https://www.nih.gov/coronavirus>.

Find NCBI SARS-CoV-2 literature, sequence, and clinical content: <https://www.ncbi.nlm.nih.gov/sars-cov-2/>.

FULL TEXT LINKS



J Dent. 2015 Aug;43(8):955-64. doi: 10.1016/j.jdent.2015.06.003. Epub 2015 Jun 22.

## A novel colourimetric technique to assess chewing function using two-coloured specimens: Validation and application

Martin Schimmel <sup>1</sup>, Panagiotis Christou <sup>2</sup>, Hideo Miyazaki <sup>3</sup>, Demetrios Halazonetis <sup>4</sup>, François R Herrmann <sup>5</sup>, Frauke Müller <sup>6</sup>

Affiliations

PMID: 26111925 DOI: [10.1016/j.jdent.2015.06.003](https://doi.org/10.1016/j.jdent.2015.06.003)

[Free article](#)

### Abstract

**Objectives:** Chewing efficiency may be evaluated using cohesive specimen, especially in elderly or dysphagic patients. The aim of this study was to evaluate three two-coloured chewing gums for a colour-mixing ability test and to validate a new purpose built software (ViewGum®).

**Methods:** Dentate participants (dentate-group) and edentulous patients with mandibular two-implant overdentures (IOD-group) were recruited. First, the dentate-group chewed three different types of two-coloured gum (gum1-gum3) for 5, 10, 20, 30 and 50 chewing cycles. Subsequently the number of chewing cycles with the highest intra- and inter-rater agreement was determined visually by applying a scale (SA) and opto-electronically (ViewGum®, Bland-Altman analysis). The ViewGum® software determines semi-automatically the variance of hue (VOH); inadequate mixing presents with larger VOH than complete mixing. Secondly, the dentate-group and the IOD-group were compared.

**Results:** The dentate-group comprised 20 participants (10 female, 30.3±6.7 years); the IOD-group 15 participants (10 female, 74.6±8.3 years). Intra-rater and inter-rater agreement (SA) was very high at 20 chewing cycles (95.00-98.75%). Gums 1-3 showed different colour-mixing characteristics as a function of chewing cycles, gum1 showed a logarithmic association; gum2 and gum3 demonstrated more linear behaviours. However, the number of chewing cycles could be predicted in all specimens from VOH (all p<0.0001, mixed linear regression models). Both analyses proved discriminative to the dental state.

**Conclusion:** ViewGum® proved to be a reliable and discriminative tool to opto-electronically assess chewing efficiency, given an elastic specimen is chewed for 20 cycles and could be recommended for the evaluation of chewing efficiency in a clinical and research setting.

**Clinical significance:** Chewing is a complex function of the oro-facial structures and the central nervous system. The application of the proposed assessments of the chewing function in geriatrics or special care dentistry could help visualising oro-functional or dental comorbidities in dysphagic patients or those suffering from protein-energy malnutrition.

**Keywords:** Chewing gum; Edentulous mouth; Mastication; Symptom assessment.

Copyright © 2015 The Authors. Published by Elsevier Ltd.. All rights reserved.

### LinkOut - more resources

#### Full Text Sources

[Bern Open Repository and Information System](#)

[ClinicalKey](#)

[Elsevier Science](#)

#### Medical

[ClinicalTrials.gov](#)